arXiv:gr-qc/0602094v1 23 Feb 2006

Some Aspects of Virtual Black Holes

Mir Faizal
IUCAA, Pune

February 6, 2008

Abstract

In this paper we analyze certain aspects of virtual black holes. We see how quantum bubble is the representative of creation and annihilation of a pair of virtual black holes that satisfy the Wheeler-DeWitt equation. We calculate the loss of quantum coherence due to these virtual black holes and also see how these virtual black holes give space-time an intrinsic entropy.

1 Introducation:-

It is expected that space-time will have quantum fluctuations in it[1]. These fluctuations are expected to dominate as we measure the space-time at smaller and smaller lengths. Finally being one at Planck length [2]. Thus space-time get a foam like structure near Planck length. This is called space-time foam[3]. There have been two models proposed to study space-time foam till now. The first model proposed to study the space-time foam was called the wormhole picture[4]. In it we took space-time to be multiple connected. So we took the space-time manifold to have a large value of first Betti number B_1. We took $B_2 = 0$ in this case [5]. However it was realized that this picture let to the wrong value of Q.C.D. θ-parameter [6]. Macroscopic wormholes also led to the wrong value of the black hole entropy [5]. So this picture was discarded and an alternative picture of space-time foam was proposed by S.W. Hawking, which was called the quantum bubble picture [5]. In it we take $B_1 = B_3 = 0$ and we also take a very large value of second Betti number B_2. Thus the space-time manifold is simple connected in this case. Quantum bubble picture predicted the correct value of Q.C.D. θ-parameter. In this picture there is also an elegant way to describe black hole evaporation. Macroscopic real black holes evaporate down to Planck size by radiating Hawking radiation. At this stage they are left with no energy or charge. Then they disappear in a sea of virtual black holes. Thus this picture seems to be a more realistic picture of space-time foam.

In this paper we will analyze some aspects of quantum bubble picture. We will start this paper by reviving the basic bubbles that are used in the construction of space-time foam.

2 Basic Quantum Bubbles

There are five basic bubbles that can be taken to be the building blocks of space-time foam i.e., $S^2 \times S^2(\chi = 4, \tau = 0), CP^2(\chi = 3, \tau = 1), \overline{CP^2}(\chi = 3, \tau = -1), K3(\chi = 24, \tau = 16), \overline{K3}(\chi = 24, \tau = -16)$. However the need of space-time to have a spin structure rules out CP^2 and $\overline{CP^2}$ and the contributions of $K3$ and $\overline{K3}$ are suppressed by Atiyah-Singer theorem as they contain Fermions zero mode[6]. So we are left with $S^2 \times S^2$ as the basic building block of space-time foam. $S^2 \times S^2$ has the symmetry group of $SO(3) \times SO(3)$ however $S^2 \times S^2 - \{p\}$ has the symmetry group of $U(1) \times U(1)$. Here p is a point on the bubble. If we take a spatial section of $S^2 \times S^2 - \{p\}$ we get $S^2 \times S^1 - \{p\}$. The topology of $S^2 \times S^1 - \{p\}$ looks like the mouth of two wormholes. We also know that spatial section of a black hole looks like a wormhole. So the topology $S^2 \times S^2 - \{p\}$ seems to look like the topology of a pair of black holes. Further we know that in Ernst solution a pair of black holes can be created by electric or magnetic fields[7]. This is similar to the pair creation in ordinary particle physics. Now as the quantum bubbles are basically quantum fluctuations in the metric, so we can interpit the bubble with topology $S^2 \times S^2 - \{p\}$ as a virtual black hole loop. Here we deform the loop on which the black hole moves into an oval with bottom representing the creation and top representing the annihilation of a pair of black holes.

3 Virtual Black Holes

We need electric or magnetic charges to produce a pair of black holes from Ernst solution. However the quantum bubbles form even in absence of any field. The reason for that is that virtual black holes are not classical geometries and hence need not satisfy Einsteins equations. They are rather solutions of the Wheeler-DeWitt equation. The solution of Wheeler-DeWitt equation is a wave function on space of geometries . Wheeler-DeWitt equation is given by [8].

$$\left[m_p G_{ijkl} \frac{\delta}{\delta h_{ij} \delta h_{kl}} - m_p h^{\frac{1}{23}} K + h^{\frac{1}{2}} T^m_m \left[\phi_o, \frac{-\delta}{\delta \phi_o}\right]\right] \psi(h_{ij}, \phi_o) = 0 \qquad (1)$$

Here m_p is the Planck mass, T^m_m is the energy momentum tensor and G_{ijkl} is given by

$$G_{ijkl} = \frac{1}{2}\sqrt{h}(h_{ik}h_{jl} + h_{il}h_{jk} - h_{ij}h_{kl}) \qquad (2)$$

They also need to satisfy the following momentum constraint equation

$$\left[-2im_p \left[\frac{\delta}{\delta h_{ij}}\right]_{|j} + T^m_m \left[\phi_o, \frac{\delta}{\delta \phi_o}\right]\right] \psi(h_{ij}, \phi_o) = 0 \qquad (3)$$

Now we can expand $\psi(h_{ij}, \phi_o)$ in terms of creation and annihilation operators.

$$\psi(h_{ij}, \phi_o) = \sum_i (\beta_{1i} a_i + \beta^\dagger_{2i} a^\dagger_i) \qquad (4)$$

And

$$\overline{\psi}(h_{ij}, \phi_o) = \sum_i (\beta_{2i} a_i + \beta^\dagger_{1i} a^\dagger_i) \tag{5}$$

Here $a^\dagger_i$ is the creation operator for a black hole and β_{1i} and β_{2i} are the complete set of bases for the conformally invariant wave equation. Now we can form a closed loop from these creation and annihilation operators. If $|0\rangle$ is the vacuum state of geometries from which we can generate the black hole geometry, and α is the amplitude for the formation of a quantum bubble, than we can write

$$\alpha = \langle 0|\overline{a^\dagger(h_{ij}, \phi_o) a(h'_{ij}, \phi'_o)} \;\; \overline{a^\dagger(h'_{ij}, \phi'_o) a(h_{ij}, \phi_o)}|0\rangle \tag{6}$$

However it is difficult to solve the Wheeler-DeWitt equation as it is faced with the problem of factor ordering. Factor ordering deals with the problem that arises due to the fact that it does not matter how we order momentum in classical equations but it does matter when they become operators. However we can get some insight about the virtual black holes from the fact that they satisfy Wheeler-DeWitt equation. The fact that they satisfy Wheeler-DeWitt equation and need not not satisfy Einstein equations means that they need not have any electric or magnetic charges that would be required by Ernst solution for pair creation of black holes. Even though they are not Classical geometries we can expect that they will have the basic feature of real black holes like an event horizon and hence the no-hair theorem will hold for them also. A particle that will fall into a the virtual black hole will lose all information except that which is allowed by the no-hair theorem[9]. So this particle will be radiated as a combination of some other particles before the annihilation of the pair of virtual black holes. This will cause loss of information to take place universally in space-time as an intrinsic property of space-time.

4 Loss of Quantum Coherence

It is well known that in case of real black holes the falling of certain states in the black hole leads to non-unitary evolution and hence loss of quantum coherence. A similar effect happens for the virtual black holes. There is an elegant formalism called the Superscattering formalism given by S.W.Hawking to describe it[5]. In it we start from taking a complete set of creation operators for states of a quantum field theory at past and future infinity. Let these set of creation operators at past and future infinity be P_i and F_i respectively. We define a Superscattering operator as one that maps a density matrix at past infinity to one at future infinity. So if ρ_- and ρ_+ are density matrix at past and future infinity respectively and \$ is the Superscattering operator, than we can write

$$\rho_+ = \$\rho_- \tag{7}$$

If the initial state is $|\psi_1\rangle\langle\psi_2|$ and the final state is $|\psi_3\rangle\langle\psi_4|$, then we have

$$\$ = \langle P^\dagger_2 F_3 F^\dagger_4 P_1\rangle \tag{8}$$

If the set of creation operators are a complete set at both past and future infinity then we have

$$\langle P^\dagger_2 F_3 F^\dagger_4 P_1\rangle = \langle P^\dagger_2 F_3\rangle\langle F^\dagger_4 P_1\rangle \tag{9}$$

Here $\langle P_2^\dagger F_3 \rangle$ is the S-matrix and $\langle F_4^\dagger P_1 \rangle$ is its adjoint. So we can write

$$\$ = SS^\dagger \tag{10}$$

Thus we can write

$$\$\rho = S\rho S^\dagger \tag{11}$$

Then using the conservation of probability we get

$$Tr(\$\rho) = 1 \text{ if } \ Tr(\rho) = 1 \tag{12}$$

Then we have

$$1 = Tr(\$\rho) = Tr(SS^\dagger \rho) = Tr(\rho) \tag{13}$$

This will hold if we have

$$SS^\dagger = 1 \tag{14}$$

This means that the S-matrix is unitary. Now to see how the density matrix evolves we have to take the trace of the future density matrix, which is given by

$$Tr(\rho_+^2) = Tr((\$\rho_-)(\$\rho_-)) \tag{15}$$

As the Supperscattering matrix factorises we can write

$$Tr((\$\rho_-)(\$\rho_-)) = Tr(S\rho_- S^\dagger S\rho_- S^\dagger) \tag{16}$$

This can be written as

$$Tr(S\rho_- S^\dagger S\rho_- S^\dagger) = Tr(S\rho_-^2 S^\dagger) = Tr(\rho_-^2) \tag{17}$$

This is the trace of the past density matrix and thus the evolution here is unitary
Thus for the case where the Superscattering matrix factorizes we have unitary evolution and hence no loss of quantum coherence. However if the creation operators do not form a complete set at past or future infinity then the Superscattering matrix will not factorize and hence the evolution will be non-unitary and so the quantum coherence will be lost. In case of real black holes F_i do not form a complete set of bases and in case of a virtual black hole both F_i and P_i do not form complete set of bases. So like real black holes, virtual black holes also lead to loss of quantum coherence. As every where in space-time we have a certain probability for the existence of virtual black holes so we can never build a truly unitary theory and quantum coherence will be lost universally.

Now we take the state inside and outside the event horizon at past infinity as $|IP_i\rangle$ and $|OP_j\rangle$ and the state inside and outside the event horizon at future infinity as $|IF_k\rangle$ and $|OF_l\rangle$ respectively. Than the state at past infinity $|P\rangle$ and future infinity $|F\rangle$ can be written as follows

$$|P\rangle = \lambda^{ij} |IP_i\rangle \times |OP_j\rangle \tag{18}$$

And

$$|F\rangle = \lambda^{kl} |IF_k\rangle \times |OF_l\rangle \tag{19}$$

Now the extent to which quantum coherence is lost at past and future infinity will be got by taking the trace over $|IP_i\rangle$ and $|IF_k\rangle$ respectively.

So if τ_P and τ_F denote the measure of loss of quantum coherence at past and future infinity than we have

$$\tau_P = 1 - Tr(\rho_-^2) = 1 - \lambda^{ij}\lambda^{ab}\overline{\lambda}_{ib}\overline{\lambda}_{aj} \tag{20}$$

And

$$\tau_F = 1 - Tr(\rho_+^2) = 1 - \lambda^{kl}\lambda^{cd}\overline{\lambda}_{kd}\overline{\lambda}_{cl} \tag{21}$$

Now if we define the total information lost as τ, than it will be given by

$$\tau = \tau_P + \tau_F \tag{22}$$

If no quantum coherence is lost than $\tau = 0$ and the Superscattering matrix will factorize. For any other value of τ the Superscattering matrix will not factorize.

To proceed further we need to first review certain topological invariants in quantum gravity[10]. Apart from the simple two well known topological invariants in quantum gravity i.e., the signature and Eulers number of the space-time manifold there is another topological invariant in quantum gravity which is called the topological invariant volume. The total action **S** for space-time manifold is given by

$$\mathbf{S} = \frac{-\Lambda}{8\pi L_p^2} Vol(M, g) \tag{23}$$

Here L_p is the Planck length and $Vol(M, g)$ is the volume of the manifold and Λ is the cosmological constant.

If we redefine the metric as follows:-

$$g_{\mu\nu} = \frac{3}{|\Lambda|}\overline{g}_{\mu\nu} \quad \text{With} \quad \Lambda = \pm 3 \tag{24}$$

Then the action becomes

$$\mathbf{S} = \frac{-\overline{g}}{8\pi L_p^2 \Lambda}\overline{v}(M, \overline{g}) \tag{25}$$

Here $\overline{v}(M, \overline{g})$ is the topological invariant volume. We can also write the partition function in terms of $\overline{v}$ as follows:-

$$Z = \sum_{\overline{v}} \rho(\overline{v}) exp\left[\frac{9}{8\pi\Lambda L_p^2}\overline{v}\right] \tag{26}$$

Here $\rho(\overline{v})$ is the measure of density of topology.

Now we can see how much quantum coherence is lost for a given topological invariant volume.

If the total flux of wave function that enters $\overline{v}$ is Γ_1 and the total flux that leaves $\overline{v}$ is Γ_2, then the total information lost in $\overline{v}$ is $\eta(\overline{v})$ and it is given by

$$\eta(\overline{v}) = \frac{\Gamma_1 - \Gamma_2}{\Gamma_1} \tag{27}$$

Now as the amplitude for formation of a virtual black hole is α, we can write

$$\eta(\overline{v}) = \tau\alpha^2\overline{v} \tag{28}$$

Now we can write Γ_2 in terms of Γ_1 as follows

$$\Gamma_2 = \Gamma_1(1 - \tau\alpha^2\overline{v}) \tag{29}$$

It is obvious that if no quantum coherence is lost than $\Gamma_1 = \Gamma_2$ as $\tau = 0$ in this case.

5 Entropy of Space-time

The fact that virtual black hole picture make the space-time to have an intrinsic entropy can be seen in various ways.
The fact that it lead to a mixed state will imply that the space-time has an entropy given by

$$S = \overline{v}\alpha^2 \left[Tr(\rho_- In\rho_-^{-1}) + Tr(\rho_+ In\rho_+^{-1})\right] \tag{30}$$

This will be zero for a pure state when no quantum coherence is lost.

Another way of looking at the intrinsic entropy of space-time is by observing that the fact that space-time gets a coarse structure at Planck length due to the presence of virtual black holes. As here different microstates will lead to the same macrostate i.e., the smooth space-time manifold at length much larger than Planck length, so space-time will have an intrinsic entropy. Here the intrinsic entropy of space-time will be given by

$$S = k\ lnW \tag{31}$$

Here W is the number of microstates generate by the virtual black holes and k is the Boltzmanns constant.

But probable the best way to look at the intrinsic entropy of space-time generate by virtual black holes is by black hole thermodynamics. It is well known that real black holes have entropy proportional to one fourth of the surface area of the event horizon [9]. If real black holes have entropy than so will the virtual black holes. This will directly imply that space-time has an intrinsic entropy.

If A is the area of a single virtual black hole than we can write the intrinsic entropy of space-time per invariant topological volume as

$$S = \frac{AN}{4L_p^2} \tag{32}$$

Here N is the number of virtual black holes per invariant topological volume $\overline{v}$. N will be equal to twice the product of probability for the formation of a virtual black hole loop and the invariant topological volume.

$$N = 2\alpha^2\overline{v} \tag{33}$$

We know that the action here is proportional to the volume of the space-time manifold. So for large volumes the action also becomes large. Thus the contributions from virtual black holes with radius larger than the Planck length will get suppressed in the partition function.
It can also be shown that the quantum fluctuations in the metric are given by [1].

$$\Delta g = -1 + \sqrt{1 + \frac{L_p^2}{L^2}} \tag{34}$$

Here Δg are the fluctuations in the metric and L is the length at which we measure the geometry. Thus at Planck length the quantum fluctuations in

geometry become of order one. So it is not possible to define geometries less than Planck length. However if we assume the radius of virtual black holes to be less than the Planck length that is what we will be implying. So the virtual black holes can not have radius less than the Planck length.
Thus the radius of the virtual black holes should be taken equal to Planck length. So we can write the area of the quantum bubble with topology $S^2 \times S^2 - \{p\}$ as follows:-

$$A = 16\pi^2 L_p^4 \tag{35}$$

Thus we can write the intrinsic entropy of space-time per topological invariant volume as follows:-

$$S = 8\pi^2 L_p^2 \alpha^2 \overline{v} = \frac{8\pi^2 L_p^2}{\tau} \eta(\overline{v}) \tag{36}$$

This is the intrinsic entropy of space-time calculated from black hole thermodynamics.
Thus it can be shown in more than one way that virtual black holes will lead space-time to have an intrinsic entropy.

6 Conclusion

Quantum bubble picture seems to be the more correct picture of space-time foam as compaired to the wormhole picture. The bubble of topology $S^2 \times S^2 - \{p\}$ is the building block of space-time and is representative of virtual black hole loop. Virtual black holes are solutions to Wheeler-DeWitt equation and hence need not have any electric or magnetic charges as would be required if they satisfied Einstein equations. Virtual black holes lead to the loss of quantum coherence, which we have calculated in this paper. Virtual black holes also lead the space-time to have an intrinsic entropy.

7 References

1. L.G.Garay - gr-qc/9911002
2. M.A.Marlov - quantum violation of topology in small spatial regions - Preprient P - 0187 of Inst. Nucl. Rev. Moscow - 1980
3. S.W.Hawking - Nucl. Phys - B114, 349 -1978
4. S.W.Hawking - Phys. Rev - D37, 904 - 1988
5. S.W.Hawking - hept-th/9510029
6. J.Preskill, S.P.Trivedi and M.B.Wise - Phys. Lett - B223, 26 - 1989
7. F.J.Ernst - J. Math. Phys - 17, 515 - 1976
8. J.B.Hartle and S.W.Hawking - Phys. Rev - D28, 2960 - 1983
9. S.W.Hawking - Commun. Math. Phys - 43, 199 - 1975
10. S.Carlip - Class. Quantum. Grav - 15, 2629 - 1998